

Polynomially presented properties: preservation proposition  
perspectives  
CUNI MFF Algebra Colloquium

Nicolas Daans

Université de Mons, Faculty of Science, Department of Mathematics, Section Mathematical Logic

24 February 2026

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

“ $-1$  is a sum of 3 squares”

“There is an element other than 0 which is both a square and minus a square”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“ $-1$  is a sum of 3 squares”

“There is an element other than 0 which is both a square and minus a square”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“−1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

We can also consider properties which an element  $X$  or tuple of elements  $(X_1, \dots, X_n)$  of the field may or may not have.

“ $X$  is a sum of 3 squares”

“ $X_1, X_2, X_3$  all have square roots”

“ $X$  is either equal to 2, or has a square root”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

We can also consider properties which an element  $X$  or tuple of elements  $(X_1, \dots, X_n)$  of the field may or may not have.

“ $X$  is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$$

“ $X_1, X_2, X_3$  all have square roots”

“ $X$  is either equal to 2, or has a square root”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

We can also consider properties which an element  $X$  or tuple of elements  $(X_1, \dots, X_n)$  of the field may or may not have.

“ $X$  is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$$

“ $X_1, X_2, X_3$  all have square roots”

$$\exists Y_1, Y_2, Y_3 : X_1 = Y_1^2, X_2 = Y_2^2, X_3 = Y_3^2$$

“ $X$  is either equal to 2, or has a square root”

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

We can also consider properties which an element  $X$  or tuple of elements  $(X_1, \dots, X_n)$  of the field may or may not have.

“ $X$  is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$$

“ $X_1, X_2, X_3$  all have square roots”

$$\exists Y_1, Y_2, Y_3 : X_1 = Y_1^2, X_2 = Y_2^2, X_3 = Y_3^2$$

“ $X$  is either equal to 2, or has a square root”

$$\exists Y : X = 2 \text{ or } X = Y^2$$

## Properties described by equations

Consider properties which a field  $K$  may or may not have:

“7 is twice the cube of an element”

$$\exists Y : 7 = 2Y^3$$

“-1 is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$$

“There is an element other than 0 which is both a square and minus a square”

~~$$\exists Y, Z_1, Z_2 : Y \neq 0, Y = Z_1^2, Y = -Z_2^2$$~~

~~$$\exists Y, Y', Z_1, Z_2 : YY' = 1, Y = Z_1^2, Y = -Z_2^2$$~~

$$\exists Y, Z_1, Z_2 : 1 = YZ_1^2, 1 = -YZ_2^2$$

We can also consider properties which an element  $X$  or tuple of elements  $(X_1, \dots, X_n)$  of the field may or may not have.

“ $X$  is a sum of 3 squares”

$$\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$$

“ $X_1, X_2, X_3$  all have square roots”

$$\exists Y_1, Y_2, Y_3 : X_1 = Y_1^2, X_2 = Y_2^2, X_3 = Y_3^2$$

“ $X$  is either equal to 2, or has a square root”

~~$$\exists Y : X = 2 \text{ or } X = Y^2$$~~

$$\exists Y : (X - 2)(X - Y^2) = 0$$

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3$

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3 \rightarrow$  needs 1 variable

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3 \rightarrow$  needs 1 variable
- $\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3 \rightarrow$  needs 1 variable
- $\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$   
 $\rightarrow$  is equivalent to  $\exists Y_1, Y_2 : -1 = Y_1^2 + Y_2^2$ , so needs only 2 variables!  
(**Proof:** If  $-1 = y_1^2 + y_2^2 + y_3^2$ , then either  $-1 = y_1^2$ , or  $-1 = \left(\frac{y_2 + y_1 y_3}{1 + y_1^2}\right)^2 + \left(\frac{y_3 - y_1 y_2}{1 + y_1^2}\right)^2$ .)

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3 \rightarrow$  needs 1 variable
- $\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$   
 $\rightarrow$  is equivalent to  $\exists Y_1, Y_2 : -1 = Y_1^2 + Y_2^2$ , so needs only 2 variables!  
(**Proof:** If  $-1 = y_1^2 + y_2^2 + y_3^2$ , then either  $-1 = y_1^2$ , or  $-1 = \left(\frac{y_2 + y_1 y_3}{1 + y_1^2}\right)^2 + \left(\frac{y_3 - y_1 y_2}{1 + y_1^2}\right)^2$ .)
- $\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$
- $\exists Y_1, Y_2, Y_3 : X_1 = Y_1^2, X_2 = Y_2^2, X_3 = Y_3^2$

## Properties described by equations

We consider properties of fields (or tuples of elements of fields) which can be described in terms of the solvability of a set of equations.

Given such a property, how many (auxiliary) variables are minimally needed to express them?

- $\exists Y : 7 = 2Y^3 \rightarrow$  needs 1 variable
- $\exists Y_1, Y_2, Y_3 : -1 = Y_1^2 + Y_2^2 + Y_3^2$   
 $\rightarrow$  is equivalent to  $\exists Y_1, Y_2 : -1 = Y_1^2 + Y_2^2$ , so needs only 2 variables!  
(**Proof:** If  $-1 = y_1^2 + y_2^2 + y_3^2$ , then either  $-1 = y_1^2$ , or  $-1 = \left(\frac{y_2 + y_1 y_3}{1 + y_1^2}\right)^2 + \left(\frac{y_3 - y_1 y_2}{1 + y_1^2}\right)^2$ .)
- $\exists Y_1, Y_2, Y_3 : X = Y_1^2 + Y_2^2 + Y_3^2$  maybe needs 3 variables ?
- $\exists Y_1, Y_2, Y_3 : X_1 = Y_1^2, X_2 = Y_2^2, X_3 = Y_3^2$  maybe needs 3 variables ?

# Existential rank

Semi-formally:

## Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

# Existential rank

Semi-formally:

## Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

The answer to the above question will be called the existential rank of  $\mathcal{P}$  (with respect to  $\mathcal{K}$ ).

# Existential rank

Semi-formally:

## Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

The answer to the above question will be called the existential rank of  $\mathcal{P}$  (with respect to  $\mathcal{K}$ ).

E.g. With respect to the class of all fields, the property “ $-1$  is a sum of 3 squares” has existential rank at most 2.

# Existential rank

Let  $\mathcal{K}$  be the class of fields of characteristic different from 2. What is the existential rank of the property “ $X_1$  and  $X_2$  are both squares”?

## Existential rank

Let  $\mathcal{K}$  be the class of fields of characteristic different from 2. What is the existential rank of the property “ $X_1$  and  $X_2$  are both squares”?

### Proposition

Let  $K$  be a field,  $\text{char}(K) \neq 2$ ,  $x_1, x_2 \in K$ . Then the following are equivalent:

- 1  $\exists y_1, y_2 \in K : x_1 = y_1^2 \text{ and } x_2 = y_2^2$ ,
- 2  $\exists z \in K : (x_1 = x_2 = 0) \text{ or } (z^4(x_1 - x_2)^2 - 2z^2(x_1 + x_2) = 1)$ .

In particular, the existential rank of “ $X_1$  and  $X_2$  are both squares” with respect to  $\mathcal{K}$  is 1.

## Existential rank

Let  $\mathcal{K}$  be the class of fields of characteristic different from 2. What is the existential rank of the property “ $X_1$  and  $X_2$  are both squares”?

### Proposition

Let  $K$  be a field,  $\text{char}(K) \neq 2$ ,  $x_1, x_2 \in K$ . Then the following are equivalent:

- ①  $\exists y_1, y_2 \in K : x_1 = y_1^2$  and  $x_2 = y_2^2$ ,
- ②  $\exists z \in K : (x_1 = x_2 = 0)$  or  $(z^4(x_1 - x_2)^2 - 2z^2(x_1 + x_2) = 1)$ .

In particular, the existential rank of “ $X_1$  and  $X_2$  are both squares” with respect to  $\mathcal{K}$  is 1.

### Proof.

(1)  $\Rightarrow$  (2): If  $x_1 \neq 0$  or  $x_2 \neq 0$ , then without loss of generality  $y_1 + y_2 \neq 0$ . Set  $z = \frac{1}{y_1 + y_2}$ .

## Existential rank

Let  $\mathcal{K}$  be the class of fields of characteristic different from 2. What is the existential rank of the property “ $X_1$  and  $X_2$  are both squares”?

### Proposition

Let  $K$  be a field,  $\text{char}(K) \neq 2$ ,  $x_1, x_2 \in K$ . Then the following are equivalent:

- 1  $\exists y_1, y_2 \in K : x_1 = y_1^2$  and  $x_2 = y_2^2$ ,
- 2  $\exists z \in K : (x_1 = x_2 = 0)$  or  $(z^4(x_1 - x_2)^2 - 2z^2(x_1 + x_2) = 1)$ .

In particular, the existential rank of “ $X_1$  and  $X_2$  are both squares” with respect to  $\mathcal{K}$  is 1.

### Proof.

- (1)  $\Rightarrow$  (2): If  $x_1 \neq 0$  or  $x_2 \neq 0$ , then without loss of generality  $y_1 + y_2 \neq 0$ . Set  $z = \frac{1}{y_1 + y_2}$ .
- (2)  $\Rightarrow$  (1): If  $x_1 \neq 0$  or  $x_2 \neq 0$ , then  $z \neq 0$ . Set  $y_1 = \frac{1+z^2(x_1-x_2)}{2z}$  and  $y_2 = \frac{1+z^2(x_2-x_1)}{2z}$ .  $\square$

# Outline

- 1 Introduction: optimally representing properties by solvability of polynomial equations  
✓
- 2 Crash course: Preservation theorems in model theory and algebra
- 3 Existential rank revisited
- 4 Outlook & a lot of open questions

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

- constant symbols (e.g. 0, 1, c, e,  $\emptyset$ , ...),

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

- constant symbols (e.g.  $0, 1, c, e, \emptyset, \dots$ ),
- function symbols (e.g.  $+, -, \cdot, \exp(\cdot), \sin(\cdot), \frac{d}{dx}, \dots$ ),

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

- constant symbols (e.g.  $0, 1, c, e, \emptyset, \dots$ ),
- function symbols (e.g.  $+, -, \cdot, \exp(\cdot), \sin(\cdot), \frac{d}{dx}, \dots$ ),
- relation symbols (e.g.  $<, \in, |, \dots$ )

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

- constant symbols (e.g.  $0, 1, c, e, \emptyset, \dots$ ),
- function symbols (e.g.  $+, -, \cdot, \exp(\cdot), \sin(\cdot), \frac{d}{dx}, \dots$ ),
- relation symbols (e.g.  $<, \in, |, \dots$ )

Function and relation symbols come with an arity (= number of inputs)

E.g.  $+, -, \cdot, <, \in$  are 2-ary (binary),  $\exp(\cdot), \sin(\cdot)$  are 1-ary (unary).

# First-order structures

Let  $\mathcal{L}$  be a first-order language, i.e. a collection of:

- constant symbols (e.g.  $0, 1, c, e, \emptyset, \dots$ ),
- function symbols (e.g.  $+, -, \cdot, \exp(\cdot), \sin(\cdot), \frac{d}{dx}, \dots$ ),
- relation symbols (e.g.  $<, \in, |, \dots$ )

Function and relation symbols come with an arity (= number of inputs)

E.g.  $+, -, \cdot, <, \in$  are 2-ary (binary),  $\exp(\cdot), \sin(\cdot)$  are 1-ary (unary).

An  $\mathcal{L}$ -structure  $\mathcal{A}$  consists of a set  $A$ , and

- for every constant symbol  $c$ , an element  $c^{\mathcal{A}} \in A$ ,
- for every  $n$ -ary function symbol  $f$ , a function  $f^{\mathcal{A}} : A^n \rightarrow A$ ,
- for every  $n$ -ary relation symbol  $R$ , a set  $R^{\mathcal{A}} \subseteq A^n$ .

# First-order structures

Main example for this presentation: the language of rings  $\mathcal{L}_{\text{ring}}$  consists of:

- two constant symbols  $0, 1$ ,
- three binary function symbols  $+, -, \cdot$ .

# First-order structures

Main example for this presentation: the language of rings  $\mathcal{L}_{\text{ring}}$  consists of:

- two constant symbols  $0, 1$ ,
- three binary function symbols  $+, -, \cdot$ .

By definition, an  $\mathcal{L}_{\text{ring}}$ -structure  $\mathcal{A}$  consists of a set  $A$  endowed with

- two elements  $0^{\mathcal{A}}, 1^{\mathcal{A}} \in A$ ,
- three binary functions  $+^{\mathcal{A}}, -^{\mathcal{A}}, \cdot^{\mathcal{A}}$ .

# First-order structures

Main example for this presentation: the language of rings  $\mathcal{L}_{\text{ring}}$  consists of:

- two constant symbols  $0, 1$ ,
- three binary function symbols  $+, -, \cdot$ .

By definition, an  $\mathcal{L}_{\text{ring}}$ -structure  $\mathcal{A}$  consists of a set  $A$  endowed with

- two elements  $0^{\mathcal{A}}, 1^{\mathcal{A}} \in A$ ,
- three binary functions  $+^{\mathcal{A}}, -^{\mathcal{A}}, \cdot^{\mathcal{A}}$ .

Note: not every  $\mathcal{L}_{\text{ring}}$ -structure is a ring.

## First-order structures

Properties which an  $\mathcal{L}$ -structure may or may not satisfy can be described and studied via first-order sentences.

## First-order structures

Properties which an  $\mathcal{L}$ -structure may or may not satisfy can be described and studied via first-order sentences.

**Example:** an  $\mathcal{L}_{\text{ring}}$ -ring structure  $\mathcal{A}$  is a ring if and only if the following sentences are satisfied:

$$\forall X, Y, Z((X + Y) + Z = X + (Y + Z) \wedge (X \cdot Y) \cdot Z = X \cdot (Y \cdot Z)),$$

$$\forall X, Y(X + Y = Y + X),$$

$$\forall X(X \cdot 1 = X \wedge 1 \cdot X = X \wedge X + 0 = X),$$

$$\forall W, X, Y, Z((W + X) \cdot (Y + Z) = ((W \cdot Y + W \cdot Z) + X \cdot Y) + X \cdot Z),$$

$$\forall X, Y, Z(X - Y = Z \leftrightarrow X = Z + Y).$$

## First-order structures

Properties which an  $\mathcal{L}$ -structure may or may not satisfy can be described and studied via first-order sentences.

**Example:** an  $\mathcal{L}_{\text{ring}}$ -ring structure  $\mathcal{A}$  is a ring if and only if the following sentences are satisfied:

$$\forall X, Y, Z((X + Y) + Z = X + (Y + Z) \wedge (X \cdot Y) \cdot Z = X \cdot (Y \cdot Z)),$$

$$\forall X, Y(X + Y = Y + X),$$

$$\forall X(X \cdot 1 = X \wedge 1 \cdot X = X \wedge X + 0 = X),$$

$$\forall W, X, Y, Z((W + X) \cdot (Y + Z) = ((W \cdot Y + W \cdot Z) + X \cdot Y) + X \cdot Z),$$

$$\forall X, Y, Z(X - Y = Z \leftrightarrow X = Z + Y).$$

The ring is a field if it additionally satisfies

$$\forall X, Y(X \cdot Y = Y \cdot X) \quad \text{and} \quad \forall X \exists Y(X = 0 \vee X \cdot Y = 1).$$

## First-order structures

Properties which an  $\mathcal{L}$ -structure may or may not satisfy can be described and studied via first-order sentences.

**Example:** an  $\mathcal{L}_{\text{ring}}$ -ring structure  $\mathcal{A}$  is a ring if and only if the following sentences are satisfied:

$$\forall X, Y, Z((X + Y) + Z = X + (Y + Z) \wedge (X \cdot Y) \cdot Z = X \cdot (Y \cdot Z)),$$

$$\forall X, Y(X + Y = Y + X),$$

$$\forall X(X \cdot 1 = X \wedge 1 \cdot X = X \wedge X + 0 = X),$$

$$\forall W, X, Y, Z((W + X) \cdot (Y + Z) = ((W \cdot Y + W \cdot Z) + X \cdot Y) + X \cdot Z),$$

$$\forall X, Y, Z(X - Y = Z \leftrightarrow X = Z + Y).$$

The ring is a field if it additionally satisfies

$$\forall X, Y(X \cdot Y = Y \cdot X) \quad \text{and} \quad \forall X \exists Y(X = 0 \vee X \cdot Y = 1).$$

A set of  $\mathcal{L}$ -sentences is called an  $\mathcal{L}$ -theory. An  $\mathcal{L}$ -structure in which all sentences of a given  $\mathcal{L}$ -theory  $T$  holds is called a model of  $T$ .

## First-order structures

Properties which an  $\mathcal{L}$ -structure may or may not satisfy can be described and studied via first-order sentences.

**Example:** an  $\mathcal{L}_{\text{ring}}$ -ring structure  $\mathcal{A}$  is a ring if and only if the following sentences are satisfied:

$$\forall X, Y, Z((X + Y) + Z = X + (Y + Z) \wedge (X \cdot Y) \cdot Z = X \cdot (Y \cdot Z)),$$

$$\forall X, Y(X + Y = Y + X),$$

$$\forall X(X \cdot 1 = X \wedge 1 \cdot X = X \wedge X + 0 = X),$$

$$\forall W, X, Y, Z((W + X) \cdot (Y + Z) = ((W \cdot Y + W \cdot Z) + X \cdot Y) + X \cdot Z),$$

$$\forall X, Y, Z(X - Y = Z \leftrightarrow X = Z + Y).$$

The ring is a field if it additionally satisfies

$$\forall X, Y(X \cdot Y = Y \cdot X) \quad \text{and} \quad \forall X \exists Y(X = 0 \vee X \cdot Y = 1).$$

A set of  $\mathcal{L}$ -sentences is called an  $\mathcal{L}$ -theory. An  $\mathcal{L}$ -structure in which all sentences of a given  $\mathcal{L}$ -theory  $T$  holds is called a model of  $T$ . E.g. An  $\mathcal{L}_{\text{ring}}$ -structure  $\mathcal{A}$  is a field if and only if it is a model of the theory  $T$  consisting of all sentences on this slide.

## $\exists$ -formulas and $\forall$ -formulas

$\mathcal{L}$ -sentences of the respective forms

$$\forall Y_1, Y_2, \dots, Y_m \psi(Y_1, \dots, Y_m) \quad \text{and} \quad \exists Y_1, \dots, Y_m \psi(Y_1, \dots, Y_m)$$

where  $\psi(Y_1, \dots, Y_m)$  is a Boolean combination of atomic formulas (in the case of  $\mathcal{L}_{\text{ring}}$ : polynomial equalities), are called universal  $\mathcal{L}$ -formulas ( $\forall$ -formulas) and existential  $\mathcal{L}$ -formulas ( $\exists$ -formulas) respectively.

A  $\forall$ -theory (respectively  $\exists$ -theory) is a theory consisting only of  $\forall$ -sentences (respectively  $\exists$ -sentences).

## $\exists$ -formulas and $\forall$ -formulas

$\mathcal{L}$ -sentences of the respective forms

$$\forall Y_1, Y_2, \dots, Y_m \psi(Y_1, \dots, Y_m) \quad \text{and} \quad \exists Y_1, \dots, Y_m \psi(Y_1, \dots, Y_m)$$

where  $\psi(Y_1, \dots, Y_m)$  is a Boolean combination of atomic formulas (in the case of  $\mathcal{L}_{\text{ring}}$ : polynomial equalities), are called universal  $\mathcal{L}$ -formulas ( $\forall$ -formulas) and existential  $\mathcal{L}$ -formulas ( $\exists$ -formulas) respectively.

A  $\forall$ -theory (respectively  $\exists$ -theory) is a theory consisting only of  $\forall$ -sentences (respectively  $\exists$ -sentences).

How does one recognize an  $\forall$ -theory ( $\exists$ -theory)?

## $\exists$ -formulas and $\forall$ -formulas

**Definition:** When  $\mathcal{A}$  is an  $\mathcal{L}$ -structure on set  $A$ , and if  $B \subseteq A$  contains all constants  $c^{\mathcal{A}}$  and is closed under all functions  $f^{\mathcal{A}}$ , then there is a natural induced  $\mathcal{L}$ -structure  $\mathcal{B}$  on  $B$  via restriction; we call  $\mathcal{B}$  an  $\mathcal{L}$ -substructure of  $\mathcal{A}$  and write  $\mathcal{B} \subseteq \mathcal{A}$ .

## $\exists$ -formulas and $\forall$ -formulas

**Definition:** When  $\mathcal{A}$  is an  $\mathcal{L}$ -structure on set  $A$ , and if  $B \subseteq A$  contains all constants  $c^{\mathcal{A}}$  and is closed under all functions  $f^{\mathcal{A}}$ , then there is a natural induced  $\mathcal{L}$ -structure  $\mathcal{B}$  on  $B$  via restriction; we call  $\mathcal{B}$  an  $\mathcal{L}$ -substructure of  $\mathcal{A}$  and write  $\mathcal{B} \subseteq \mathcal{A}$ .

### Proposition

*Every  $\mathcal{L}_{\text{ring}}$ -substructure of a ring is a ring.*

### Proof.

All  $\mathcal{L}_{\text{ring}}$ -sentences (axioms) for rings are  $\forall$ -sentences. When they hold in a structure, then also in every substructure. □

## $\exists$ -formulas and $\forall$ -formulas

**Definition:** When  $\mathcal{A}$  is an  $\mathcal{L}$ -structure on set  $A$ , and if  $B \subseteq A$  contains all constants  $c^{\mathcal{A}}$  and is closed under all functions  $f^{\mathcal{A}}$ , then there is a natural induced  $\mathcal{L}$ -structure  $\mathcal{B}$  on  $B$  via restriction; we call  $\mathcal{B}$  an  $\mathcal{L}$ -substructure of  $\mathcal{A}$  and write  $\mathcal{B} \subseteq \mathcal{A}$ .

### Proposition

*Every  $\mathcal{L}_{\text{ring}}$ -substructure of a ring is a ring.*

### Proof.

All  $\mathcal{L}_{\text{ring}}$ -sentences (axioms) for rings are  $\forall$ -sentences. When they hold in a structure, then also in every substructure. □

### Proposition

*The class of  $\mathcal{L}_{\text{ring}}$ -structures which are fields can not be described by an  $\forall$ -theory.*

### Proof.

$\mathbb{Q}$  is a field, but its  $\mathcal{L}_{\text{ring}}$ -substructure  $\mathbb{Z}$  is not. □

## $\exists$ -formulas and $\forall$ -formulas

### Theorem (Łoś-Tarski Preservation Theorem ( $\forall$ -version), 1954)

*Let  $T$  be an  $\mathcal{L}$ -theory. Suppose that every substructure of a model of  $T$  is again a model of  $T$ . Then  $T$  is equivalent to an  $\forall$ -theory.*

*(Call two theories equivalent if they have the same models.)*

## $\exists$ -formulas and $\forall$ -formulas

### Theorem (Łoś-Tarski Preservation Theorem ( $\forall$ -version), 1954)

*Let  $T$  be an  $\mathcal{L}$ -theory. Suppose that every substructure of a model of  $T$  is again a model of  $T$ . Then  $T$  is equivalent to an  $\forall$ -theory.*

*(Call two theories equivalent if they have the same models.)*

Dually: one has

### Theorem (Łoś-Tarski Preservation Theorem ( $\exists$ -version), 1954)

*Let  $T$  be an  $\mathcal{L}$ -theory. Suppose that every substructure of a non-model of  $T$  is again a non-model of  $T$ . Then  $T$  is equivalent to an  $\exists$ -theory.*

## $\exists$ -formulas and $\forall$ -formulas

**Example:** Add to the theory of rings the  $\mathcal{L}$ -sentence

$$\exists X_1, X_2, X_3, X_4 \forall Y (X_1 = Y \vee X_2 = Y \vee X_3 = Y \vee X_4 = Y).$$

This axiom expresses precisely that the ring has at most 4 elements.

## $\exists$ -formulas and $\forall$ -formulas

**Example:** Add to the theory of rings the  $\mathcal{L}$ -sentence

$$\exists X_1, X_2, X_3, X_4 \forall Y (X_1 = Y \vee X_2 = Y \vee X_3 = Y \vee X_4 = Y).$$

This axiom expresses precisely that the ring has at most 4 elements.

Clearly, a subring of a ring with at most 4 elements also has at most 4 elements. Hence, by the Preservation Theorem, the sentence above must be equivalent to an  $\forall$ -sentence.

## $\exists$ -formulas and $\forall$ -formulas

**Example:** Add to the theory of rings the  $\mathcal{L}$ -sentence

$$\exists X_1, X_2, X_3, X_4 \forall Y (X_1 = Y \vee X_2 = Y \vee X_3 = Y \vee X_4 = Y).$$

This axiom expresses precisely that the ring has at most 4 elements.

Clearly, a subring of a ring with at most 4 elements also has at most 4 elements. Hence, by the Preservation Theorem, the sentence above must be equivalent to an  $\forall$ -sentence.

For example (by Pigeonhole Principle):

$$\forall Y_1, Y_2, Y_3, Y_4, Y_5 \left( \bigvee_{1 \leq i < j \leq 5} (Y_i = Y_j) \right)$$

## Another example: $\forall\exists$ -formulas

$\mathcal{L}$ -sentences of the form

$$\forall Y_1, \dots, Y_m \exists Z_1, \dots, Z_n \psi(Y_1, \dots, Y_m, Z_1, \dots, Z_n)$$

where  $\psi$  is a Boolean combination of atomic formulas, are called universal-existential  $\mathcal{L}$ -formulas ( $\forall\exists$ -formulas).

An  $\forall\exists$ -theory is a theory consisting of only  $\forall\exists$ -sentences.

## Another example: $\forall\exists$ -formulas

$\mathcal{L}$ -sentences of the form

$$\forall Y_1, \dots, Y_m \exists Z_1, \dots, Z_n \psi(Y_1, \dots, Y_m, Z_1, \dots, Z_n)$$

where  $\psi$  is a Boolean combination of atomic formulas, are called universal-existential  $\mathcal{L}$ -formulas ( $\forall\exists$ -formulas).

An  $\forall\exists$ -theory is a theory consisting of only  $\forall\exists$ -sentences.

How does one recognize an  $\forall\exists$ -theory?

## Another example: $\forall\exists$ -formulas

**Definition:** When  $\mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$  is an ascending chain of  $\mathcal{L}$ -structures with underlying sets  $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$ , the union  $\bigcup_{i \in \mathbb{N}} \mathcal{A}_i$  naturally carries an  $\mathcal{L}$ -structure, denoted  $\bigcup_{i \in \mathbb{N}} \mathcal{A}_i$  and called the directed union.

## Another example: $\forall\exists$ -formulas

**Definition:** When  $\mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$  is an ascending chain of  $\mathcal{L}$ -structures with underlying sets  $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$ , the union  $\bigcup_{i \in \mathbb{N}} A_i$  naturally carries an  $\mathcal{L}$ -structure, denoted  $\bigcup_{i \in \mathbb{N}} \mathcal{A}_i$  and called the directed union.

### Proposition

*Let  $K_0 \subseteq K_1 \subseteq \dots$  be an ascending chain of fields. Then  $\bigcup_{i \in \mathbb{N}} K_i$  is also a field.*

### Proof.

All  $\mathcal{L}_{\text{ring}}$ -sentences (axioms) for fields are  $\forall\exists$ -sentences. When they hold in an ascending chain of structures, then also in their union. □

## Another example: $\forall\exists$ -formulas

**Definition:** When  $\mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$  is an ascending chain of  $\mathcal{L}$ -structures with underlying sets  $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$ , the union  $\bigcup_{i \in \mathbb{N}} A_i$  naturally carries an  $\mathcal{L}$ -structure, denoted  $\bigcup_{i \in \mathbb{N}} \mathcal{A}_i$  and called the directed union.

### Proposition

*Let  $K_0 \subseteq K_1 \subseteq \dots$  be an ascending chain of fields. Then  $\bigcup_{i \in \mathbb{N}} K_i$  is also a field.*

### Proof.

All  $\mathcal{L}_{\text{ring}}$ -sentences (axioms) for fields are  $\forall\exists$ -sentences. When they hold in an ascending chain of structures, then also in their union. □

### Theorem (Chang-Łoś-Suszko Preservation Theorem, 1960's)

*Let  $T$  be an  $\mathcal{L}$ -theory. Suppose that, whenever  $\mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \mathcal{A}_2 \subseteq \dots$  is a chain of models of  $T$ , also  $\bigcup_{i \in \mathbb{N}} \mathcal{A}_i$  is a model of  $T$ . Then  $T$  is equivalent to an  $\forall\exists$ -theory.*

# Outline

- ① Introduction: optimally representing properties by solvability of polynomial equations ✓
- ② Crash course: Preservation theorems in model theory and algebra ✓
- ③ Existential rank revisited
- ④ Outlook & a lot of open questions

## Recall: existential rank

Recall the original question:

### Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

## Recall: existential rank

Recall the original question:

### Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

The answer to the above question will be called the existential rank of  $\mathcal{P}$  (with respect to  $\mathcal{K}$ ).

## Recall: existential rank

Recall the original question:

### Question

*Let  $\mathcal{K}$  be a class of fields,  $n \in \mathbb{N}$ . Let  $\mathcal{P}(X_1, \dots, X_n)$  be a property which a  $n$ -tuple of elements  $(x_1, \dots, x_n)$  in a field  $K \in \mathcal{K}$  may or may not have.*

*What is the smallest  $m \in \mathbb{N}$  (if it exists) for which the property  $\mathcal{P}(X_1, \dots, X_n)$  is equivalent to the solvability of  $S(X_1, \dots, X_n, Y_1, \dots, Y_m)$  in  $K$  (for  $Y_1, \dots, Y_m$ ) for all  $K \in \mathcal{K}$ , where  $S$  is some boolean combination of polynomial equalities?*

The answer to the above question will be called the existential rank of  $\mathcal{P}$  (with respect to  $\mathcal{K}$ ).

We are looking for a criterion for a property  $\mathcal{P}$  to be given as

$$\exists Y_1, \dots, Y_m \psi$$

where  $\psi$  is a Boolean combination of polynomial equations.

# The property “ $X_1, \dots, X_n$ are squares”

Consider the property  $\pi_n(X_1, \dots, X_n)$  stating that  $X_1, \dots, X_n$  are all squares, i.e. given by

$$\exists Y_1, \dots, Y_n (X_1 = Y_1^2 \wedge \dots \wedge X_n = Y_n^2).$$

# The property “ $X_1, \dots, X_n$ are squares”

Consider the property  $\pi_n(X_1, \dots, X_n)$  stating that  $X_1, \dots, X_n$  are all squares, i.e. given by

$$\exists Y_1, \dots, Y_n (X_1 = Y_1^2 \wedge \dots \wedge X_n = Y_n^2).$$

## Proposition

*Let  $\mathcal{K}$  be the class of all fields,  $n \in \mathbb{N}$ . Then  $\pi_n$  has existential rank  $n$  with respect to  $\mathcal{K}$ .*

*Proof on blackboard.*

# The property “ $X_1, \dots, X_n$ are squares”

For contrast, we have:

## Proposition

*Let  $\mathcal{K}$  be the class of all fields  $K$  with  $\text{char}(K) \neq 2$ , let  $n \geq 1$ . Then  $\pi_n$  has existential rank 1 with respect to  $\mathcal{K}$ .*

# The property “ $X_1, \dots, X_n$ are squares”

For contrast, we have:

## Proposition

Let  $\mathcal{K}$  be the class of all fields  $K$  with  $\text{char}(K) \neq 2$ , let  $n \geq 1$ . Then  $\pi_n$  has existential rank 1 with respect to  $\mathcal{K}$ .

## Proof idea.

- 1 If  $K$  is any field with  $\text{char}(K) \neq 2$  and  $a_1, \dots, a_n \in K$ , the extension  $K(\sqrt{a_1}, \dots, \sqrt{a_n})/K$  is generated by a single element by the *Primitive Element Theorem*.



# The property “ $X_1, \dots, X_n$ are squares”

For contrast, we have:

## Proposition

Let  $\mathcal{K}$  be the class of all fields  $K$  with  $\text{char}(K) \neq 2$ , let  $n \geq 1$ . Then  $\pi_n$  has existential rank 1 with respect to  $\mathcal{K}$ .

## Proof idea.

- 1 If  $K$  is any field with  $\text{char}(K) \neq 2$  and  $a_1, \dots, a_n \in K$ , the extension  $K(\sqrt{a_1}, \dots, \sqrt{a_n})/K$  is generated by a single element by the *Primitive Element Theorem*.
- 2 A quantitative version of the Łoś-Tarski Preservation Theorem (D., Dittmann, Fehm, 2021) relates existential rank to the minimal number of generators of extensions.



# The property “ $X_1, \dots, X_n$ are squares”

For contrast, we have:

## Proposition

Let  $\mathcal{K}$  be the class of all fields  $K$  with  $\text{char}(K) \neq 2$ , let  $n \geq 1$ . Then  $\pi_n$  has existential rank 1 with respect to  $\mathcal{K}$ .

## Proof idea.

- 1 If  $K$  is any field with  $\text{char}(K) \neq 2$  and  $a_1, \dots, a_n \in K$ , the extension  $K(\sqrt{a_1}, \dots, \sqrt{a_n})/K$  is generated by a single element by the *Primitive Element Theorem*.
- 2 A quantitative version of the Łoś-Tarski Preservation Theorem (D., Dittmann, Fehm, 2021) relates existential rank to the minimal number of generators of extensions.



*Note:* It is possible to find an explicit formula with 1 existential quantifier for all  $n$  (Becher, D., ca. 2019, only in my PhD thesis).

# Existential rank and essential dimension

In “Existential rank and essential dimension of diophantine sets” (D., Dittmann, Fehm, 2021) we develop systematically the connection between two related concepts of complexity:

- Existential rank of properties in fields,

# Existential rank and essential dimension

In “Existential rank and essential dimension of diophantine sets” (D., Dittmann, Fehm, 2021) we develop systematically the connection between two related concepts of complexity:

- Existential rank of properties in fields,
- essential and canonical dimension of varieties (as introduced by Merkurjev, Berhuy, Favi, 2003 - based on earlier work of Buhler and Reichstein, 1997).

Informally: the essential dimension of an algebraic object is the minimal number of algebraically independent parameters needed to describe it.

## Existential rank and essential dimension

For example, by using results of Karpenko and Merkurjev (2003) which imply that the projective quadratic over  $\mathbb{Q}(T)$  given by the equation

$$TX_0^2 = X_1^2 + \dots + X_n^2$$

has canonical dimension  $n - 1$ , we obtain:

### Proposition

*Let  $\mathcal{K}$  be the class of all fields of characteristic 0,  $n \in \mathbb{N}$ . Then the formula*

$$\exists Y_1, \dots, Y_n : X = Y_1^2 + \dots + Y_n^2$$

*has existential rank  $n$ .*

# Outline

- 1 Introduction: optimally representing properties by solvability of polynomial equations ✓
- 2 Crash course: Preservation theorems in model theory and algebra ✓
- 3 Existential rank revisited ✓
- 4 Outlook & a lot of open questions

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ?

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,
  - ⑥ (Pasten, 2021) there exists properties over  $\mathbb{C}(T)$  of arbitrarily large existential rank,

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2(\!(T)\!)$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,
  - ⑥ (Pasten, 2021) there exists properties over  $\mathbb{C}(T)$  of arbitrarily large existential rank,
  - ⑦  $K = \mathbb{Q}$ : the property “ $X$  is a sum of two squares” has existential rank 2.

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,
  - ⑥ (Pasten, 2021) there exists properties over  $\mathbb{C}(T)$  of arbitrarily large existential rank,
  - ⑦  $K = \mathbb{Q}$ : the property “ $X$  is a sum of two squares” has existential rank 2.
- Things we do not know include:

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,
  - ⑥ (Pasten, 2021) there exists properties over  $\mathbb{C}(T)$  of arbitrarily large existential rank,
  - ⑦  $K = \mathbb{Q}$ : the property “ $X$  is a sum of two squares” has existential rank 2.
- Things we do not know include:
  - ①  $K = \mathbb{Q}$ : Does there exist a property of existential rank greater than 2?

# Outlook

- So far: consider existential rank with respect to class  $\mathcal{K}$  of *all fields* (or all extensions of a certain base field).  
There is a clear model-theoretic/functorial/geometric perspective.
- What if  $\mathcal{K}$  consists of a *single field*  $K$ ? Examples of things we know:
  - ①  $K = \mathbb{C}$  or  $K$  finite: every existential property has existential rank 0,
  - ②  $K = \mathbb{R}$  or  $K = \mathbb{Q}_p$ : every existential property has existential rank at most 1,
  - ③  $K = \mathbb{F}_2((T))$ : the property “ $X_1, \dots, X_n$  are all squares” has existential rank  $n$ ,
  - ④ BUT if  $K/\mathbb{F}_2$  is finitely generated, then the property “ $X_1, \dots, X_n$  are all squares” has existential rank 1,
  - ⑤ there exists  $K$  for which the property “ $X$  is a sum of  $n$  squares” has existential rank  $n$ ,
  - ⑥ (Pasten, 2021) there exists properties over  $\mathbb{C}(T)$  of arbitrarily large existential rank,
  - ⑦  $K = \mathbb{Q}$ : the property “ $X$  is a sum of two squares” has existential rank 2.
- Things we do not know include:
  - ①  $K = \mathbb{Q}$ : Does there exist a property of existential rank greater than 2?
  - ② If  $K$  has a property of existential rank larger than 1, does it have properties of arbitrarily large existential rank?

# Thanks for your attention!

- [BF03] Grégory Berhuy and Giordano Favi. “Essential Dimension: A functorial point of view (after A. Merkurjev)”. In: [Documenta Mathematica](#) 8 (2003), pp. 279–330.
- [Daa22] Nicolas Daans. “Existential first-order definitions and quadratic forms”. Available as [hdl.handle.net/10067/1903760151162165141](https://hdl.handle.net/10067/1903760151162165141). PhD thesis. University of Antwerp, 2022.
- [DDF21] Nicolas Daans, Philip Dittmann, and Arno Fehm. “Existential rank and essential dimension of diophantine sets”. Available as [arXiv:2102.06941](https://arxiv.org/abs/2102.06941). 2021.
- [KM03] Nikita Karpenko and Alexander Merkurjev. “Essential dimension of quadrics”. In: [Inventiones Mathematicae](#) 153.2 (2003), pp. 361–372.
- [Pas22] Hector Pasten. “Notes on the DPRM property for listable structures”. In: [The Journal of Symbolic Logic](#) 87.1 (2022), pp. 273–312.

Nicolas Daans

E-mail: [nicolas.daans@umons.ac.be](mailto:nicolas.daans@umons.ac.be)